

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/05, 38/06, C07K 5/00	A1	(11) International Publication Number: WO 99/17790 (43) International Publication Date: 15 April 1999 (15.04.99)
(21) International Application Number: PCT/US98/21055 (22) International Filing Date: 7 October 1998 (07.10.98) (30) Priority Data: 60/061,309 7 October 1997 (07.10.97) US 09/166,808 6 October 1998 (06.10.98) US (71) Applicant: CEPHALON INC. [US/US]; 145 Brandywine Parkway, West Chester, PA 19380 (US). (72) Inventors: CHATTERJEE, Sankar; 228 Henley Road, Wynnewood, PA 19096 (US). MALLAMO, John, P.; 616 Font Road, Glenmoore, PA 19343 (US). BIHOVSKY, Ron; 804 Primrose Lane, Wynnewood, PA 19096 (US). WELLS, Gregory, J.; 818 Serpentine Drive, West Chester, PA 19382 (US). (74) Agents: MILLER, Suzanne, E. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris LLP, 46th floor, One Liberty Place, Philadelphia, PA 19103 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PEPTIDE-CONTAINING α -KETOAMIDE CYSTEINE AND SERINE PROTEASE INHIBITORS (57) Abstract This invention relates to peptide-containing α -ketoamide inhibitors of cysteine and serine proteases, methods for making these compounds, and methods for using the same.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

PEPTIDE-CONTAINING α -KETOAMIDE CYSTEINE AND SERINE PROTEASE
INHIBITORS

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims benefit of U.S.

- 5 Provisional Application Serial No. 60/061,309, filed October 7, 1997, and U.S. Application entitled "Peptide-containing α -Ketoamide Cysteine and Serine Protease Inhibitors," filed October 6, 1998, the disclosures of which are hereby incorporated herein by reference in their entirety.

10 FIELD OF THE INVENTION

This invention relates to peptide-containing α -ketoamide inhibitors of cysteine and serine proteases, methods for making these compounds, and methods for using the same.

15 BACKGROUND OF THE INVENTION

- Numerous cysteine and serine proteases have been identified in human tissues. A "protease" is an enzyme which degrades proteins into smaller components (peptides). The terms "cysteine protease" and "serine protease" refer
20 to proteases which are distinguished by the presence therein of a cysteine or serine residue which plays a

- 2 -

critical role in the catalytic process. Mammalian systems, including humans, normally degrade and process proteins via a variety of enzymes including cysteine and serine proteases. However, when present at elevated levels or
5 when abnormally activated, cysteine and serine proteases may be involved in pathophysiological processes.

For example, calcium-activated neutral proteases ("calpains") comprise a family of intracellular cysteine proteases which are ubiquitously expressed in mammalian
10 tissues. Two major calpains have been identified; calpain I and calpain II. While calpain II is the predominant form in many tissues, calpain I is thought to be the predominant form in pathological conditions of nerve tissues. The calpain family of cysteine proteases has been implicated in
15 many diseases and disorders, including neurodegeneration, stroke, Alzheimer's, amyotrophy, motor neuron damage, acute central nervous system injury, muscular dystrophy, bone resorption, platelet aggregation, cataracts and inflammation. Calpain I has been implicated in excitatory
20 amino-acid induced neurotoxicity disorders including ischemia, hypoglycemia, Huntington's Disease, and epilepsy. The lysosomal cysteine protease cathepsin B has been implicated in the following disorders: arthritis, inflammation, myocardial infarction, tumor metastasis, and
25 muscular dystrophy. Other lysosomal cysteine proteases include cathepsins C, H, L and S. Interleukin-1 β converting enzyme ("ICE") is a cysteine protease which catalyzes the formation of interleukin-1 β . Interleukin-1 β is an immunoregulatory protein implicated in the following
30 disorders: inflammation, diabetes, septic shock, rheumatoid arthritis, and Alzheimer's disease. ICE has also been linked to apoptotic cell death of neurons, which is implicated in a variety of neurodegenerative disorders including Parkinson's disease, ischemia, and amyotrophic
35 lateral sclerosis (ALS).

- 3 -

Cysteine proteases are also produced by various pathogens. The cysteine protease clostripain is produced by *Clostridium histolyticum*. Other proteases are produced by *Trypanosoma cruzi*, malaria parasites *Plasmodium*

- 5 *falciparum* and *P. vinckei* and *Streptococcus*. Hepatitis A viral protease HAV C3 is a cysteine protease essential for processing of picornavirus structural proteins and enzymes.


- Exemplary serine proteases implicated in degenerative disorders include thrombin, human leukocyte
10 elastase, pancreatic elastase, chymase and cathepsin G. Specifically, thrombin is produced in the blood coagulation cascade, cleaves fibrinogen to form fibrin and activates Factor VIII; thrombin is implicated in thrombophlebitis, thrombosis and asthma. Human leukocyte elastase is
15 implicated in tissue degenerative disorders such as rheumatoid arthritis, osteoarthritis, atherosclerosis, bronchitis, cystic fibrosis, and emphysema. Pancreatic elastase is implicated in pancreatitis. Chymase, an enzyme important in angiotensin synthesis, is implicated in
20 hypertension, myocardial infarction, and coronary heart disease. Cathepsin G is implicated in abnormal connective tissue degradation, particularly in the lung.

- Given the link between cysteine and serine proteases and various debilitating disorders, compounds which inhibit
25 these proteases would be useful and would provide an advance in both research and clinical medicine. The present invention is directed to these, as well as other, important ends.

- 4 -

SUMMARY OF THE INVENTION

The present invention is directed to selected peptide-containing α -ketoamide inhibitors of cysteine and serine proteases represented by the general formula I:



I

wherein:

Q has the formula $G-B-(CHR^4)_v$, where R^4 is independently H or alkyl having from 1 to 4 carbons;

10 v is 0, 1, or 2;

B is selected from the group consisting of $C(=O)$, $OC(=O)$, $S(=O)_m$, CH_2 , a bond, $NR^5C(=O)$, $S(=O)_m-A-C(=O)$, and $C(=O)-A-C(=O)$, where R^5 is H or lower alkyl;

m is 0, 1, or 2;

15 A is lower alkylene or cycloalkylene, optionally substituted with one or more halogen atoms, aryl, or heteroaryl groups;

M is a carbon atom;

G is selected from the group consisting of H, a
 20 blocking group, lower alkyl, lower alkenyl, aryl having from about 6 to about 14 carbons, heterocyclyl having from about 5 to about 14 ring atoms, heterocycloalkyl having from about 5 to about 14 ring atoms, arylalkyl having from about 7 to about 15 carbons, heteroarylalkyl, and arylheteroalkyl
 25 wherein the aryl portion can be unfused or fused with the heteroalkyl ring, said alkyl, aryl, heterocyclyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, and arylheteroalkyl groups being optionally substituted with one or more J groups;

30 J is selected from the group consisting of halogen, CN, nitro, lower alkyl, cycloalkyl, heterocycloalkyl, heteroalkyl, halogenated alkyl, aryloxyalkyl, alkylthio,

- 5 -

alkylsulfonyl, aryl, heteroaryl, arylalkyl, arylalkyloxy, arylsulfonyl, heteroarylsulfonyl, alkoxycarbonyl, alkoxyalkyl, acyl, alkoxy, hydroxy, carboxy, hydroxyalkyl, amino, alkylamino, and aminoalkyl, said amino group or said
5 amino group of said aminoalkyl or alkylamino group being optionally substituted with an acyl group, an alkoxy group, or with 1 to 3 aryl, lower alkyl, cycloalkyl, or alkoxyalkyl groups; and said aryl, heteroaryl, heterocycloalkyl, and heteroalkyl groups being further optionally substituted by a
10 J group;

each Aaa is independently an amino acid which optionally contains one or more blocking groups;

n is 0, 1, 2, or 3;

R¹ and R² are independently selected from the group
15 consisting of H, alkyl having from one to about 6 carbons, arylalkyl having from about 7 to about 15 carbons, heteroalkyl in which the ring contains from about 5 to about 14 ring atoms, heteroarylalkyl in which the heteroaryl ring contains from about 5 to about 14 ring atoms, alkoxyalkyl, a
20 side chain of a naturally occurring amino acid in the R or S configuration, and (CH₂)_pNH-L, said alkyl, arylalkyl, heteroalkyl, heteroarylalkyl, and alkoxyalkyl groups being optionally substituted with one or more J groups;

p is 0, 1, 2, or 3;

25 L is selected from the group consisting of alkoxycarbonyl having from 2 to about 7 carbons, arylalkoxycarbonyl in which the arylalkoxy group contains about 7 to about 15 carbons, and S(=O)₂R^ε;

R⁶ is selected from the group consisting of
30 lower alkyl, and aryl having from about 6 to about 14 carbons;

R³ is selected from the group consisting of H, alkyl having from one to about 6 carbons, arylalkyl having from about 7 to about 15 carbons, heteroalkyl in which the ring
35 contains from about 5 to about 14 ring atoms,


- 6 -

heteroarylalkyl in which the heteroaryl ring contains from about 5 to about 14 ring atoms, alkoxyalkyl, a side chain of a naturally occurring amino acid in the R or S configuration, $(CH_2)_pNH-L$, $C(=O)R^7$, $S(=O)_2R^7$, a blocking group, and when combined with the carbon atom to which R^1 is attached an alkylene group having from 2 to 5 carbons, said alkylene group being optionally substituted with a group selected from the group consisting of aryl, azide, CN, a protected amino group, and OSO_2 -aryl, said alkyl, arylalkyl, heteroalkyl, heteroarylalkyl, and alkoxyalkyl groups being optionally substituted with one or more J groups;

R^7 is selected from the group consisting of aryl having from about 6 to about 14 carbons, heteroaryl having from about 5 to about 14 ring atoms, arylalkyl having from about 7 to about 15 carbons, alkyl having from 1 to about 10 carbons, said aryl, heteroaryl, arylalkyl and alkyl groups being optionally substituted with one or more J groups, heteroalkyl having from 2 to about 7 carbons, alkoxy having from about 1 to about 10 carbons, and amino optionally substituted with 1 or more alkyl groups;

q is 0 or 1;


Z is selected from the group consisting of $C(=O)C(=O)NH-X-A^1-K$ and



25 X is a bond or -O-;

A^1 is the same as A;

K is selected from the group consisting of $N(R^{10})Y$,



- 7 -

D is a fused aryl or heteroaryl group;

R¹¹ is selected from the group consisting of alkoxy, aryloxy, and NHR¹²;

R¹² is selected from the group consisting of H, alkyl, aryl, and heteroaryl, said alkyl, aryl or heteroaryl groups being optionally substituted with one or more J groups;

Y is selected from the group consisting of SO₂R³, C(=O)NHR⁹, C(=S)NHR⁹, C(=NCN)R¹¹, C(=NC(=O)NHR¹⁰)R¹¹, and CO₂R⁸;

R⁸ is selected from the group consisting of alkyl, alkoxy, aryl, and heterocyclyl, said alkyl, alkoxy, aryl, or heterocyclyl groups being optionally substituted with one or more J groups;

R⁵ is selected from the group consisting of H, alkyl, aryl, and heteroaryl, said alkyl, aryl, or heteroaryl groups being optionally substituted with one or more J groups;

or an R⁹ alkyl group may be combined with an A¹ alkylene group to form a N-containing heterocyclic 5- or 6-membered ring;

R¹⁰ is selected from the group consisting of H and lower alkyl;

or in the moiety SO₂N(R⁵)R¹⁰, R⁵ and R¹⁰ may be combined together with the N atom to which they are attached to form a N-containing heterocyclic 5- or 6- membered ring;

or where A¹ is an alkylene group, and K is N(R¹⁰)Y wherein R¹⁰ is alkyl, said R¹⁰ alkyl group may be combined with said A¹ alkylene group to form a N-containing heterocyclic 5- or 6- membered ring; or a pharmaceutically acceptable salt thereof.

In some preferred embodiments of the compounds of Formula I, n and v are each 0, q is 1, B is a bond, and G is H. In further preferred embodiments of the compounds of Formula I, R¹ is the sidechain of a naturally occurring amino acid. In still further preferred embodiments of the compounds of Formula I, R³ is -S(=O)₂R⁷.

- 8 -

In some preferred embodiments of the compounds of Formula I, R^2 is benzyl or alkoxyalkyl. In more preferred embodiments, X is a bond, and Y is SO_2R^8 . Preferably, A^1 is $-CH_2-CH_2-$, $-CH_2-CH(CH_3)-$, or $-(CH_3)CH-CH_2-$.

5 In further preferred embodiments of the compounds of Formula I, R^1 is a serine sidechain, which is optionally capped with a benzyl group. Preferably, the carbon to which the serine sidechain is attached, designated "M" in Formula I, is a carbon atom in the D configuration.

10 In preferred embodiments of the compounds of Formula I, R^2 is benzyl, R^7 is methyl, and R^8 is substituted phenyl, unsubstituted phenyl, substituted heteroaryl, or unsubstituted heteroaryl. In particularly preferred
15 embodiments, R^8 is aryl, aryl substituted with amino, aryl substituted with heterocyclomethyl, heteroaryl, alkyl substituted with heteroaryl, or heteroaryl substituted with alkylthio, haloalkyl, alkyl, phenylsulfonyl, halogen, aminophenyl, amino, or dialkylaminoalkyl.

In further preferred embodiments of the compounds of
20 Formula I, n, v and q are each 0, B is $(C=O)$, and G is phenyl or lower alkyl, said phenyl or lower alkyl groups being optionally substituted with one or more J groups.

In more preferred embodiments of the invention, n and v are each 0, q is 1, R^1 is the side chain of an amino acid in
25 the D- or L-configuration, R^3 is $S(=O)_2R^7$, G is H, B is a bond, R^2 is benzyl or alkoxyalkyl, X is a bond, and Y is SO_2R^8 .

In other preferred embodiments, A^1 is CH_2CH_2 , $CH_2CH(CH_3)$, or $(CH_3)CHCH_2$. In more preferred embodiments, R^1 is a serine
30 side chain in the D-configuration in which the hydroxyl group is capped with benzyl, R^2 is benzyl, R^7 is methyl, and R^8 is substituted or unsubstituted phenyl or substituted or unsubstituted heteroaryl.

More preferred are the substituents shown for R_1-R_4 , B,
35 G, Aaa, X, A^1 , Y, n, q and v shown for the compounds in

- 9 -

Tables 2, 3, 4 and 5. Especially preferred are the substituents shown for compounds 9, 13, 17, 22, and 29-151.

Some especially preferred embodiments of the compounds of Formula I are shown in Tables 2, 3, 4 and 5, *infra*, with compounds 9, 13, 17, 22, and 29-151 being particularly preferred.

Because the peptide-containing α -ketoamides of the invention inhibit cysteine proteases and serine proteases, they can be used in both research and therapeutic settings.

10 In a research environment, preferred compounds having defined attributes can be used to screen for natural and synthetic compounds which evidence similar characteristics in inhibiting protease activity. The compounds can also be used in the refinement of *in vitro* and *in vivo* models for
15 determining the effects of inhibition of particular proteases on particular cell types or biological conditions.

In a therapeutic setting, given the connection between cysteine proteases and certain defined disorders, and serine proteases and certain defined disorders, compounds of the
20 invention can be utilized to alleviate, mediate, reduce and/or prevent disorders which are associated with abnormal and/or aberrant activity of cysteine proteases and/or serine proteases.

In preferred embodiments, compositions are provided for
25 inhibiting a serine protease or a cysteine protease comprising a compound of the invention and a pharmaceutically acceptable carrier. In other preferred embodiments, methods are provided for inhibiting serine proteases or cysteine proteases comprising contacting a
30 protease selected from the group consisting of serine proteases and cysteine proteases with an inhibitory amount of a compound of the invention.


Methodologies for making the present peptide-containing α -ketoamide inhibitors are also disclosed. Other
35 useful methodologies will be apparent to those skilled in

- 10 -

the art, once armed with the present disclosure. These and other features of the compounds of the subject invention are set forth in more detail below.

DETAILED DESCRIPTION

5 Disclosed herein are the selected peptide-containing α -ketoamides which are represented by the following formula I:



I

wherein:

10 Q has the formula $\text{G}-\text{B}-(\text{CHR}^4)_v$, where R^4 is independently H or alkyl having from 1 to 4 carbons;

v is 0, 1, or 2;

B is selected from the group consisting of $\text{C}(=\text{O})$, $\text{OC}(=\text{O})$, $\text{S}(=\text{O})_m$, CH_2 , a bond, $\text{NR}^5\text{C}(=\text{O})$, $\text{S}(=\text{O})_m-\text{A}-\text{C}(=\text{O})$, and
 15 $\text{C}(=\text{O})-\text{A}-\text{C}(=\text{O})$, where R^5 is H or lower alkyl;

m is 0, 1, or 2;

A is lower alkylene or cycloalkylene, optionally substituted with one or more halogen atoms, aryl, or heteroaryl groups;

20 M is a carbon atom;

G is selected from the group consisting of H, a blocking group, lower alkyl, lower alkenyl, aryl having from about 6 to about 14 carbons, heterocyclyl having from about 5 to about 14 ring atoms, heterocycloalkyl having from about 5 to about 14 ring atoms, arylalkyl having from about 7 to
 25 about 15 carbons, heteroarylalkyl, and arylheteroalkyl wherein the aryl portion can be unfused or fused with the heteroalkyl ring, said alkyl, aryl, heterocyclyl, heterocycloalkyl, arylalkyl, heteroarylalkyl, and

- 11 -

arylheteroalkyl groups being optionally substituted with one or more J groups;

J is selected from the group consisting of halogen, CN, nitro, lower alkyl, cycloalkyl, heterocycloalkyl, heteroalkyl, halogenated alkyl, aryloxyalkyl, alkylthio, alkylsulfonyl, aryl, heteroaryl, arylalkyl, arylalkyloxy, arylsulfonyl, heteroarylsulfonyl, alkoxycarbonyl, alkoxyalkyl, acyl, alkoxy, hydroxy, carboxy, hydroxyalkyl, amino, alkylamino, and aminoalkyl, said amino group or said amino group of said aminoalkyl or alkylamino group being optionally substituted with an acyl group, an alkoxy group, or with 1 to 3 aryl, lower alkyl, cycloalkyl, or alkoxyalkyl groups; and said aryl, heteroaryl, heterocycloalkyl, and heteroalkyl groups being further optionally substituted by a J group;

each Aaa is independently an amino acid which optionally contains one or more blocking groups;

n is 0, 1, 2, or 3;

R¹ and R² are independently selected from the group consisting of H, alkyl having from one to about 6 carbons, arylalkyl having from about 7 to about 15 carbons, heteroalkyl in which the ring contains from about 5 to about 14 ring atoms, heteroarylalkyl in which the heteroaryl ring contains from about 5 to about 14 ring atoms, alkoxyalkyl, a side chain of a naturally occurring amino acid in the R or S configuration, and (CH₂)_pNH-L, said alkyl, arylalkyl, heteroalkyl, heteroarylalkyl, and alkoxyalkyl groups being optionally substituted with one or more J groups;

p is 0, 1, 2, or 3;

L is selected from the group consisting of alkoxycarbonyl having from 2 to about 7 carbons, arylalkoxycarbonyl in which the arylalkoxy group contains about 7 to about 15 carbons, and S(=O)₂R^ε;

- 12 -


R^6 is selected from the group consisting of lower alkyl, and aryl having from about 6 to about 14 carbons;

R^3 is selected from the group consisting of H, alkyl
 5 having from one to about 6 carbons, arylalkyl having from about 7 to about 15 carbons, heteroalkyl in which the ring contains from about 5 to about 14 ring atoms, heteroarylalkyl in which the heteroaryl ring contains from about 5 to about 14 ring atoms, alkoxyalkyl, a side chain of
 10 a naturally occurring amino acid in the R or S configuration, $(CH_2)_pNH-L$, $C(=O)R^7$, $S(=O)_2R^7$, a blocking group, and when combined with the carbon atom to which R^1 is attached an alkylene group having from 2 to 5 carbons, said alkylene group being optionally substituted with a group
 15 selected from the group consisting of aryl, azide, CN, a protected amino group, and OSO_2 -aryl, said alkyl, arylalkyl, heteroalkyl, heteroarylalkyl, and alkoxyalkyl groups being optionally substituted with one or more J groups;

R^7 is selected from the group consisting of aryl
 20 having from about 6 to about 14 carbons, heteroaryl having from about 5 to about 14 ring atoms, arylalkyl having from about 7 to about 15 carbons, alkyl having from 1 to about 10 carbons, said aryl, heteroaryl, arylalkyl and alkyl groups being optionally substituted with one or more J groups,
 25 heteroalkyl having from 2 to about 7 carbons, alkoxy having from about 1 to about 10 carbons, and amino optionally substituted with 1 or more alkyl groups;

q is 0 or 1;

Z is selected from the group consisting of
 30 $C(=O)C(=O)NH-X-A^1-K$ and




- 13 -

X is a bond or -O-;

A¹ is the same as A;

K is selected from the group consisting of N(R¹⁰)Y,



5 D is a fused aryl or heteroaryl group;

R¹¹ is selected from the group consisting of alkoxy, aryloxy, and NHR¹²;

R¹² is selected from the group consisting of H, alkyl, aryl, and heteroaryl, said alkyl, aryl or heteroaryl groups being optionally substituted with one or more J groups;

Y is selected from the group consisting of SO₂R⁸, C(=O)NHR⁹, C(=S)NHR⁹, C(=NCN)R¹¹, C(=NC(=O)NHR¹⁰)R¹¹, and CO₂R⁸;

R⁸ is selected from the group consisting of alkyl, alkoxy, aryl, and heterocyclyl, said alkyl, alkoxy, aryl, or heterocyclyl groups being optionally substituted with one or more J groups;

R⁹ is selected from the group consisting of H, alkyl, aryl, and heteroaryl, said alkyl, aryl, or heteroaryl groups being optionally substituted with one or more J groups;

or an R⁹ alkyl group may be combined with an A¹ alkylene group to form a N-containing heterocyclic 5- or 6-membered ring;

R¹⁰ is selected from the group consisting of H and lower alkyl;

or in the moiety SO₂N(R⁸)R¹⁰, R⁸ and R¹⁰ may be combined together with the N atom to which they are attached to form a N-containing heterocyclic 5- or 6- membered ring;

or where A¹ is an alkylene group, and K is N(R¹⁰)Y wherein R¹⁰ is alkyl, said R¹⁰ alkyl group may be combined

- 14 -

with said A¹ alkylene group to form a N-containing heterocyclic 5- or 6- membered ring; or a pharmaceutically acceptable salt thereof.

It is recognized that various stereoisomeric forms of the compounds of Formula I may exist. Preferred compounds of the invention have any Aaa groups being α -amino acids in the L-configuration. However, racemates and individual enantiomers and mixtures thereof form part of the present invention.

10 The carbon atom designated as "M" in the compounds of Formula I can exist in either the D or the L configuration. In some preferred embodiments, M is a carbon atom having the "D" configuration.

As used herein, the term "alkyl" includes straight-
15 chain, and branched hydrocarbon groups such as, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, 1-ethylpentyl, hexyl, and octyl groups. "Cycloalkyl" groups are cyclic alkyl groups, such as, for example, cyclopropyl, methylcyclopentyl, and
20 cyclohexyl groups. Preferred alkyl groups have 1 to about 10 carbon atoms, most preferably "lower alkyl" of 1 to about 6 carbon atoms. "Alkylene" groups are alkyl groups having two points of attachment; i.e., non-terminal alkyl groups. "Lower alkylene" groups are branched or unbranched alkylene
25 groups of 1 to about 6 carbon atoms such as, for example, ethylene(-CH₂CH₂-), propylene, butylene, hexylene, 1-methylethylene, 2-methylethylene, and 2-methylpropylene. "Cycloalkylene" groups are cyclic alkylene groups. "Acyl" groups are alkylcarbonyl groups. "Aryl" groups are aromatic
30 cyclic compounds preferably including but not limited to phenyl, tolyl, naphthyl, anthracyl, phenanthryl, and pyrenyl. Also included within the definition of "aryl" are ring systems having two aromatic rings connected by a bond, such as biphenyl. Preferred aryl groups include phenyl and
35 naphthyl.

- 15 -

The term "carbocyclic", as used herein, refers to cyclic groups in which the ring portion is composed solely of carbon atoms. The term "halogen" refers to F, Cl, Br, and I atoms. The term "arylalkyl" denotes alkyl groups which bear aryl groups, for example, benzyl groups. As used herein, "alkoxy" groups are alkyl groups linked through an oxygen atom. Examples of alkoxy groups include methoxy ($-OCH_3$) and ethoxy ($-OCH_2CH_3$) groups. In general, the term "oxy" when used as a suffix denotes attachment through an oxygen atom. Thus, alkoxycarbonyl groups are carbonyl groups which contain an alkoxy substituent, i.e., groups of general formula $-C(=O)-O-R$, where R is alkyl. The term "alkoxyalkyl" denotes an alkoxy group attached to an alkyl group. The term "aryloxy" denotes an aryl group linked through an oxygen atom, and the term "arylalkyloxy" denotes an arylalkyl group linked through an oxygen atom.


The terms "heterocycle", "heterocyclyl", and "heterocyclic" refer to cyclic groups in which a ring portion includes at least one heteroatom such as O, N or S. Heterocyclic groups include "heteroaryl" as well as "heteroalkyl" groups. The term "heteroaryl" denotes aryl groups having one or more hetero atoms (e.g., O, N, or S) contained within an aromatic ring. Also included within the definition of "heteroaryl" are ring systems having two aromatic rings connected by a bond, where at least one of the rings contains a hetero atom. Preferred "heteroaryl" groups include pyridyl, pyrimidyl, pyrrolyl, furyl, thienyl, imidazolyl, triazolyl, tetrazolyl, quinolyl, isoquinolyl, benzoimidazolyl, thiazolyl, bipyridyl, pyridylthiophenyl, pyrimidylthiophenyl, benzimidazolyl, isoxazolylthiophenyl, pyrazolylthiophenyl, phthalimido, and benzothiazolyl. The term "heterocycloalkyl" denotes a heterocycle attached through a lower alkyl group. The term "heteroarylalkyl" denotes a heteroaryl group attached through an alkyl group. As used herein, the term "heteroalkyl" denotes a

- 16 -













heterocyclic group which contains at least one saturated carbon atom in a heterocyclic ring. Examples of heteroalkyl groups include piperidine, dihydropyridine, and tetrahydroisoquinyl groups. The term "arylheteroalkyl" as
 5 used herein denotes a "heteroalkyl" group connected through an aryl group. One preferred example of an arylheteroalkyl group is dibenzo-γ-pyranyl.

As used herein, the term "amino acid" denotes a molecule containing both an amino group and a carboxyl
 10 group. As used herein the term "L-amino acid" denotes an α-amino acid having the L- configuration around the α-carbon, that is, a carboxylic acid of general formula $\text{CH}(\text{COOH})(\text{NH}_2)$ - (side chain), having the L-configuration. The term "D-amino acid" similarly denotes a carboxylic acid of general formula
 15 $\text{CH}(\text{COOH})(\text{NH}_2)$ - (side chain), having the D-configuration around the α-carbon. Side chains of L-amino acids include naturally occurring and non-naturally occurring moieties. Nonnaturally occurring (i.e., unnatural) amino acid side chains are moieties that are used in place of naturally
 20 occurring amino acid sidechains in, for example, amino acid analogs. See, for example, Lehninger, *Biochemistry*, Second Edition, Worth Publishers, Inc, 1975, pages 73-75. One representative amino acid side chain is the lysyl side chain, $-(\text{CH}_2)_4-\text{NH}_2$. Other representative α-amino acid side
 25 chains are shown below in Table 1.

Table 1

CH_3-	$\text{HS}-\text{CH}_2-$
$\text{HO}-\text{CH}_2-$	$\text{HO}_2\text{C}-\text{CH}(\text{NH}_2)-\text{CH}_2-\text{S}-\text{S}-\text{CH}_2-$
$\text{C}_6\text{H}_5-\text{CH}_2-$	CH_3-CH_2-
30 $\text{HO}-\text{C}_6\text{H}_4-\text{CH}_2-$	$\text{CH}_3-\text{S}-\text{CH}_2-\text{CH}_2-$
	$\text{CH}_3-\text{CH}_2-\text{S}-\text{CH}_2-\text{CH}_2-$
	$\text{HO}-\text{CH}_2-\text{CH}_2-$
	$\text{CH}_3-\text{CH}(\text{OH})-$
	$\text{HO}_2\text{C}-\text{CH}_2-\text{NHC}(=\text{O})-\text{CH}_2-$

- 17 -



Functional groups present in the compounds of Formula I may contain blocking groups. Blocking groups are known *per se* as chemical functional groups that can be selectively appended to functionalities, such as hydroxyl groups, amino groups, thio groups, and carboxyl groups. Protecting groups are blocking groups which can be readily removed from functionalities. These groups are present in a chemical compound to render such functionality inert to chemical reaction conditions to which the compound is exposed. Any

5 of a variety of protecting groups may be employed with the present invention. Examples of such protecting groups are the benzyloxycarbonyl (Cbz; Z), toluenesulfonyl, t-butoxycarbonyl, methyl ester, and benzyl ether groups.

Other preferred protecting groups according to the invention

15 may be found in Greene, T.W. and Wuts, P.G.M., "Protective

- 18 -

Groups in Organic Synthesis" 2d. Ed., Wiley & Sons, 1991, which is hereby incorporated by reference in its entirety.

Further blocking groups useful in the compounds of the present invention include those that bear acyl, aroyl, alkyl, alkanesulfonyl, arylalkanesulfonyl, or arylsulfonyl substituents on their amino groups. Other useful blocking groups include alkyl ethers, e.g., the methyl ether of serine.

The disclosed compounds of the invention are useful for the inhibition of cysteine proteases and serine proteases. As used herein, the terms "inhibit" and "inhibition" mean having an adverse effect on enzymatic activity. An inhibitory amount is an amount of a compound of the invention effective to inhibit a cysteine and/or serine protease.

Pharmaceutically acceptable salts of the cysteine and serine protease inhibitors also fall within the scope of the compounds as disclosed herein. The term "pharmaceutically acceptable salts" as used herein means an inorganic acid addition salt such as hydrochloride, sulfate, and phosphate, or an organic acid addition salt such as acetate, maleate, fumarate, tartrate, and citrate. Examples of pharmaceutically acceptable metal salts are alkali metal salts such as sodium salt and potassium salt, alkaline earth metal salts such as magnesium salt and calcium salt, aluminum salt, and zinc salt. Examples of pharmaceutically acceptable organic amine addition salts are salts with morpholine and piperidine. Examples of pharmaceutically acceptable amino acid addition salts are salts with lysine, glycine, and phenylalanine.

Compounds provided herein can be formulated into pharmaceutical compositions by admixture with pharmaceutically acceptable nontoxic excipients and carriers. As noted above, such compositions may be prepared for use in parenteral administration, particularly

- 19 -

in the form of liquid solutions or suspensions; or oral administration, particularly in the form of tablets or capsules; or intranasally, particularly in the form of powders, nasal drops, or aerosols; or dermally, via, for example, transdermal patches; or prepared in other suitable fashions for these and other forms of administration as will be apparent to those skilled in the art.

The composition may conveniently be administered in unit dosage form and may be prepared by any of the methods well known in the pharmaceutical art, for example, as described in *Remington's Pharmaceutical Sciences* (Mack Pub. Co., Easton, PA, 1980). Formulations for parenteral administration may contain as common excipients sterile water or saline, polyalkylene glycols such as polyethylene glycol, oils and vegetable origin, hydrogenated naphthalenes and the like. In particular, biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be useful excipients to control the release of the active compounds. Other potentially useful parenteral delivery systems for these active compounds include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, cyclodextrins and liposomes. Formulations for inhalation administration contain as excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or oily solutions for administration in the form of nasal drops, or as a gel to be applied intranasally. Formulations for parenteral administration may also include glycocholate for buccal administration, a salicylate for rectal administration, or citric acid for vaginal administration. Formulations for transdermal patches are preferably lipophilic emulsions.

The materials for this invention can be employed as the sole active agent in a pharmaceutical or can be used

- 20 -

in combination with other active ingredients which could facilitate inhibition of cysteine and serine proteases in diseases or disorders.

The concentrations of the compounds described herein in a therapeutic composition will vary depending upon a number of factors, including the dosage of the drug to be administered, the chemical characteristics (e.g., hydrophobicity) of the compounds employed, and the route of administration. In general terms, the compounds of this invention may be provided in effective inhibitory amounts in an aqueous physiological buffer solution containing about 0.1 to 10% w/v compound for parenteral administration. Typical dose ranges are from about 1µg/kg to about 1 g/kg of body weight per day; a preferred dose range is from about 0.01 mg/kg to 100 mg/kg of body weight per day. Such formulations typically provide inhibitory amounts of the compound of the invention. The preferred dosage of drug to be administered is likely, however, to depend on such variables as the type or extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, and formulation of the compound excipient, and its route of administration.

As used herein, the term "contacting" means directly or indirectly causing at least two moieties to come into physical association with each other. Contacting thus includes physical acts such as placing the moieties together in a container, or administering moieties to a patient. Thus, for example administering a compound of the invention to a human patient evidencing a disease or disorder associated with abnormal and/or aberrant activity of such proteases falls within the scope of the definition of the term "contacting".

The invention is further illustrated by way of the following examples which are intended to elucidate the

- 21 -

invention. These examples are not intended, nor are they to be construed, as limiting the scope of the disclosure.

Examples

General Methods. Thin layer chromatography was performed
5 on silica gel plates (MK6F 60A, size 1 x 3 in, layer
thickness 250 μm , Whatman Inc.). Preparative thin layer
chromatography was performed on silica gel plates (size 20
x 20 in, layer thickness 1000 micron, Analtech).
Preparative column chromatography was carried out using
10 Merck silica gel, 40-63 μm , 230-400 mesh. ^1H NMR spectra
were recorded on a GE QE Plus instrument (300 MHz) using
tetramethylsilane as internal standard. Electrospray mass
spectra were recorded on a VG platform II instrument
(Fisons Instruments).


15 Compounds of the invention were prepared
following one of the General Methods A, B, C or D.

Example 1

Preparation of Compound 9 by General Method A

- 22 -

General Method A



Preparation of Compound 1

This compound and related hydroxy acids used in this study were synthesized following a general procedure of Harbeson et al, *J. Med. Chem.* 1994, 37, 2918-2929, which is incorporated herein by reference in its entirety.

Preparation of Compound 2

To a cooled (-10 °C) solution of compound 1 (4.30 g, 0.015 mol) in anhydrous methanol (50 mL) was added slowly thionyl chloride (3.20 mL). After 0.5 hour, the cooling bath was removed, the mixture was stirred for an additional 16 hours and concentrated to give a residue which on trituration with ethyl acetate (30 mL) gave a white solid. The solid was separated by filtration and

- 23 -

dried to give 3.50 g of compound 2 which was used directly in the next step; MS m/e 210(M+H).

Preparation of Compound 3

The preparation of this compound is shown in
5 General Method E.

Preparation of Compound 4

To a cooled (0 °C) solution of compound 1 (1.00 g, 0.0034 mol) in anhydrous DMF (20 mL) was added NMM (1.40 g, 0.014 mmol) followed by 1-HOBt (0.54 g, 0.0040 mmol) and BOP (1.80 g, 0.0040 mmol). The mixture was stirred for 15 minutes and to it was added compound 3 (0.75g, 0.0032 mmol). The cooling bath was removed and the mixture was stirred for 4 hours, poured into ice-water (200 mL), and
15 extracted into ethyl acetate (3 x 100 mL). The organic layer was washed with 2% citric acid solution (2 x 50 mL), 2% sodium bicarbonate solution (2 x 50 mL), brine (1 x 50 mL), and it was dried over anhydrous sodium sulfate. Solvent evaporation under reduced pressure gave a crude
20 solid which was washed several times with n-pentane to produce 1.30 g of compound 4.

Compound 4: white solid (diastereomeric mixture); $^1\text{H-NMR}$ (DMSO- d_6) δ 7.90 and 7.65 (2 sets of t, 1H), 7.75 (d, 2H), 7.55 (q, 2H), 7.15 (m, 6H), 6.55 and 5.80 (2 sets of d, 1H), 3.90 (m, 2H), 3.30 (d, 1H), 3.10 (m, 2H), 2.75 (m, 2H), 2.50 (m, 3H), 1.20 (s, 9H). MS m/e 478 (M+H), 500 (M+Na).

Preparation of Compound 5

To a solution of compound 4 (0.40 g, 0.84 mmol) in 1,4-dioxane (15 mL) was added 4 N HCl in dioxane (15 mL). The reaction mixture was stirred at room temperature
30

- 24 -

for 2 hours, then concentrated at reduced pressure to give a residue which was washed several times with ethyl acetate and dried under vacuum to give 0.30 g of compound 5; $^1\text{H-NMR}$ (DMSO-d_6) showed complete absence of tBoc peak at δ 1.20 ppm; MS m/e 378 ($\text{M}+\text{H}$). This material was used directly in the next step.

Preparation of Compound 7

To a mixture of D-Ser(Bn) (compound 6, 1.00g, 5 mmol) and 1 N NaOH (10 mL, 10 mmol) at 0 °C was slowly added methanesulfonyl chloride (0.80 g, 7.69 mmol). After 0.5 hour, the cooling bath was removed, the mixture was stirred overnight and acidified (pH ~ 2-3) with 2 N HCl. The aqueous layer was extracted into ethyl acetate (3 x 50 mL). The combined organic layer was washed with water (1 x 20 mL) and brine (1 x 20 mL), and dried over MgSO_4 . Solvent evaporation gave a residue which was redissolved in methylene chloride (10 mL); addition of hexanes produced a white solid which was filtered and dried to give 1.02 g of compound 7.

Preparation of Compound 8

This compound was prepared by coupling compound 7 and compound 5, using NMM/HOBt/BOP as coupling agents, following the procedure described above for the preparation of compound 4. In some of the related examples, EDCI/HOBt were used as coupling agents.

Preparation of Compound 9

To a cooled (0 °C) solution of compound 8 (0.31 g, 0.49 mmol) in anhydrous methylene chloride (10 mL) was added Dess-Martin periodinane reagent (0.425 g, 1.00 mmol). The cooling bath was removed and the mixture was stirred for an additional 1 hour. The solution was then diluted

- 25 -


with methylene chloride (10 mL), and washed with 10% sodium thiosulfate solution (5 x 5 mL), saturated sodium bicarbonate solution (2 x 5 mL), and brine (1 x 5 mL), and dried over anhydrous sodium sulfate. Solvent removal under reduced pressure gave a residue which was washed with n-pentane (10 mL) and dried under vacuum to produce 0.178 g of compound 9; $^1\text{H-NMR}$ spectrum revealed a minor amount of epimerization had taken place.

Compound 9: white solid; $^1\text{H-NMR}$ (DMSO-d_6) δ 8.75 (t, 1H), 8.60 and 8.50 (2 doublets, 1H), 7.75 (d, 2H), 7.65-7.00 (a series of m, 15H), 5.25 (broad m, 1H), 4.45 and 4.235 (2 singlets, 2H), 4.15 (m, 1H), 3.35-2.60 (a series of m, 8H), 3.35 and 3.25 (2 singlets, 3H) MS m/e 631 (M+H), 653 (M+Na).

Example 2

Preparation of Compound 13 by General Method B

General Method B



In General Method B, compound 4, prepared as described above, was oxidized by Dess-Martin periodinane

- 26 -

- reagent to generate compound 10 which on tBoc-deprotection (2 N HCl in dioxane) produced the amine-salt, compound 11. Coupling (NMM/HOBt/BOP) of compound 11 with N-phenylsulfonyl-(L)-Pro (compound 12) yielded compound 13.
- 5 Purification was achieved by passing a solution of the crude material in methylene chloride through Sep-Pak® Vac 6cc (1 g) silica cartridge (Waters Corporation, Milford, MA), eluting with methylene chloride, followed by various combinations of methylene chloride and ethyl acetate.
- 10 Harbeson et al. (*J. Med. Chem.* 1994, 37, 2918-2929) reported that silica gel chromatography of a ketoamide epimerizes the chiral center at P₁.

Compound 13: white solid; ¹H-NMR (CDCl₃) δ 7.90-7.00 (a series of m, 18H), 5.40 and 5.30 (2 multiplets, 1H), 4.10 (m, 1H), 3.50-3.00 (m, 8H), 1.90-1.40 (m, 4H). MS m/e 613(M+H), 635(M+Na).

15


Example 3

Preparation of Compound 17 by General Method C

- In General Method C, compound 2 was coupled (NMM/HOBt/BOP) with L-Cbz-Leu to give compound 14 which was hydrolyzed (aq. NaOH) to compound 15. Coupling (NMM/HOBt/BOP) of compound 15 with compound 3 gave compound 16 which underwent Dess-Martin oxidation to generate compound 17.
- 20

- 27 -

General Method C



Compound 17: white solid; $^1\text{H-NMR}$ (CDCl_3) δ 7.85 (d, 2H), 7.60-7.00 (a series of m, 15H), 6.60 (d, 1H), 5.40 (m, 1H), 5.20 (q, 1H), 5.10 (s, 2H), 4.10 (broad, 1H), 3.50-3.00 (a series of m, 6H), 1.65-1.30 (m, 3H), 0.90 (d, 6H). MS m/e 623(M+H), 645(M+Na).

Example 4


Preparation of Compound 22 by General Method D

In General Method D, compound 7 was coupled (NMM/HOBt/BOP) with compound 2 to generate compound 18 which underwent Dess-Martin oxidation to generate compound 19. Hydrolysis (LiOH , $\text{MeOH-H}_2\text{O}$) of compound 19 gave compound 20 which was coupled (NMM/HOBt/BOP) with compound 21 to give compound 22. Compound 22 was purified by silica gel chromatography.

Compound 22: white solid; MS m/e 646(M+H), 668(M+Na).

- 28 -

General Method D




Preparation of Intermediates

Example 5

The preparation of a representative example of an amine (compound 3), containing a terminal sulfonamide moiety, is shown in General Method E.

- 29 -

General Method E**Preparation of Compound 24**

To a solution of 1,2-ethylenediamine (compound 23, 10.80 g, 12.00 mL, 0.18 mol) in THF (30 mL) was added
5 slowly BOC-ON (22.10 g, 0.09 mol) in THF (70 mL) over a period of 4 hours. The reaction mixture was stirred overnight, concentrated on a rotavapor, and taken up into water (150 mL). The aqueous layer was acidified (pH ~ 5-6) with solid citric acid monohydrate, washed with ether (3 x
10 50 mL) and then treated (at 0 °C) with 6 N NaOH solution to make it basic (pH ~ 12-13). The basic solution was extracted into ethyl acetate (3 x 100 mL), and the combined ethyl acetate layer was dried (MgSO_4) and concentrated to generate 7.23 g of monoprotected diamine, compound 24.

15 Compound 24: viscous liquid; $^1\text{H-NMR}$ (CDCl_3) δ 5.00 (broad, 1H), 3.20 (broad q, 2H), 2.80 (t, 2H), 1.45 (s, 9H), 1.25 (broad, 2H).

Preparation of Compound 25

A cooled (0-5 °C) solution of compound 24 (0.321
20 g, 0.002 mol) in methylene chloride (5 mL) was treated sequentially with triethylamine (0.243 g, 0.33 mL, 0.0024 mol) and benzenesulfonyl chloride (0.423 g, 0.30 mL, 0.0024 mol). The ice-bath was removed and the mixture was stirred

- 30 -

for an additional 0.5 hour, washed successively with water (2 x 5 mL), cold (0-5 °C) 0.5 N HCl (1 x 5 mL), 2% NaHCO₃ solution (1 x 5 mL), and brine (1 x 5 mL). The solution was dried (MgSO₄) and the solvent was evaporated to give a
5 residue which was washed several times with n-pentane. A total of 0.60 g of the sulfonamide derivative, compound 25, was obtained.

Compound 25: white solid, mp 92-95 °C; R_f (TLC, 5% methanol in methylene chloride) 0.55; ¹H-NMR (CDCl₃) δ 7.85 (d,
10 2H), 7.55 (m, 3H), 5.30 (broad d, 1H), 4.85 (broad, 1H), 3.25 (broad q, 2H), 3.10 (broad q, 2H), 1.40 (s, 9H).

Preparation of Compound 3

A solution of compound 25 (0.560 g, 0.0019 mol) in 1,4-dioxane (4 mL) was treated with 4 N HCl in dioxane
15 (4 mL). The mixture was stirred at room temperature for 1 hour and concentrated at the rotavapor. The residue was washed several times with ethyl acetate and dried under vacuum to give 0.40 g of compound 3.


Compound 3: white solid, mp 178-180 °C; ¹H-NMR (DMSO-d₆) δ
20 8.20-8.00 (broad t, 4H), 7.80 (d, 2H), 7.60 (m, 3H), 2.95 (broad q, 2H), 2.80 (broad, 2H).

Example 6

Preparation of Compound 28

The preparation of a representative example of an
25 intermediate amine (compound 28) containing a terminal biaryl sulfonamide moiety is shown in General Method F.

- 31 -

General Method F

A mixture of compound 26 (prepared from compound 24 and 5-bromothiophene-2-sulfonyl chloride, following the same general procedure as described above for the preparation of compound 25, 0.50 g, 1 eqv), dimethoxyethane (10 mL), 2 M Na₂CO₃ (5 eqv), phenylboronic acid (1.40 eqv) and Pd(PPh₃)₄ (0.04 eqv) was heated at 135 °C for 2.5 hours. The reaction mixture was concentrated at the rotavapor, and the residue was taken up into water (20 mL). The aqueous layer was acidified with citric acid and extracted into methylene chloride (3 x 20 mL). The combined organic layer was washed with water (1 x 10 mL) and brine (1 x 10 mL). It was dried (MgSO₄) and concentrated to a small volume.


Trituration of the residue with hexanes gave a solid which was separated by filtration and dried under vacuum to produce 0.37 g of compound 27; ¹H-NMR (CDCl₃) δ 7.60-7.20 (a series of m, 7H), 5.35 (broad, 1H), 4.85 (broad, 1H), 3.30 (m, 2H), 3.20 (m, 2H), 1.40 (s, 9H). For a general description of this reaction procedure, see Miyaura et al., Chem. Rev. 1995, 95, 2457-2483.

- 32 -

Compound 27 was converted to compound 28 following the procedure described for the preparation of compound 3.

Example 7**5 Preparation of Taurine Sulfonamide Intermediate**

The preparation of a representative taurine sulfonamide intermediate is shown in General Method G.

General Method G

The phthalimide of taurine, prepared by a known
10 procedure (R. Winterbottom et al., *J. Amer. Chem. Soc.*,
1947, 69, 1393 - 1401) was converted to its sulfonyl
chloride with phosphorous pentachloride in refluxing
benzene. This was allowed to react with aniline in the
presence of pyridine to form the corresponding sulfonamide.
15 The phthalimide protecting group was then removed by
refluxing with hydrazine and the resulting taurine
sulfonamide was isolated as its hydrochloride.

- 33 -

Example 8

Syntheses of compounds 29 through 50 in Tables 2 and 3 were carried out using the designated general methods, as described, and the appropriate starting materials.

Example 9**Inhibition of Cysteine Protease Activity**

To evaluate inhibitory activity, stock solutions (40 times concentrated) of each compound to be tested were prepared in 100% anhydrous DMSO and 5 μ l of each inhibitor preparation were aliquoted into each of three wells of a 96-well plate. Recombinant human calpain I, prepared by the method of Meyer et al. (*Biochem. J.* 1996, 314: 511-519; incorporated herein by reference in its entirety), was diluted into assay buffer (i.e., 50mM Tris, 50mM NaCl, 1mM EDTA, 1mM EGTA, and 5mM β -mercaptoethanol, pH 7.5, including 0.2mM Succ-Leu-Tyr-MNA), and 175 μ l was aliquoted into the same wells containing the independent inhibitor stocks as well as to positive control wells containing 5 μ l DMSO, but no compound. To start the reaction, 20 μ l of 50 mM CaCl_2 in assay buffer was added to all wells of the plate, excepting three, which were used as background signal baseline controls. Substrate hydrolysis was monitored every 5 minutes for a total of 30 minutes. Substrate hydrolysis in the absence of inhibitor was linear for up to 15 minutes.

Inhibition of calpain I activity was calculated as the percent decrease in the rate of substrate hydrolysis in the presence of inhibitor relative to the rate in its absence. Comparison between the inhibited and control rates was made within the linear range for substrate hydrolysis. The IC_{50} s of inhibitors (concentration yielding 50% inhibition) were determined from the percent decrease in rates of substrate hydrolysis in the presence of five to

- 34 -

seven different concentrations of the test compound. The results were plotted as percent inhibition versus log inhibitor concentration, and the IC_{50} was calculated by fitting the data to the four-parameter logistic equation shown below using the program GraphPad Prism (GraphPad Software, Inc., San Diego, CA.).


$$y = d + [(a-d) / (1 + (x / c)^b)]$$





The parameters a, b, c, and d are defined as follows: a is % inhibition in the absence of inhibitor, b is the slope, c is the IC_{50} , and d is the % inhibition at an infinite concentration of inhibitor.

Results are presented in Tables 2, 3, 4 and 5 below.










- 35 -

Table 2
Inhibitory Activity of Linear α -Ketoamides






5	Cmp.	W	R ₂	R	Calpain	Prep.	MS
	No.				IC ₅₀ nM	Method	(M+1) ⁺
	9	Ms-D-Ser (Bn)	Bn	Ph	16	A	631
	13	PhSO ₂ -L-Pro	Bn	Ph	(78%) **	B	613
	17	Cbz-L-Leu	Bn	Ph	11	C	623
	22	Ms-D-Ser (Bn)	Bn		42	D	646
10	29	Ms-D-Ser (Bn)	Bn		14	A	714
	30	PhSO ₂ -L-Phe	Bn	Ph	(97%) **	B	663
	31	Ms-D-Ser (Bn)	Bn		29	A	632
	32	Ms-D-Ser (Bn)	Bn		10	A	704

- 36 -

	33	Ms-D-Ser (Bn)	Bn		17	A	761
	34	Ms-D-Ser (Bn)	Bn		25	A	786 (M+2) ⁺
	35	Ms-D-Ser (Bn)	Bn	CH ₃	91	A	569
	36*	Ms-D-Ser (Bn)	CH ₂ OMe		14	A	668
5	37	Ms-D-Ser (Bn)	Bn		18	C	777
	38*	Ms-D-Ser (Bn)	CH ₂ OMe	Ph	(100%) *	A	585
	39	Ms-D-Ser (Bn)	Bn		22	C	715, 717 ⁷⁹ Br, ⁸¹ Br
	40	Ms-D-Ser (Bn)	Bn		63	D	722
	41	Ms-D-Ser (Bn)	Bn		(88%) **	D	699
10	42	Ms-D-Ser (Bn)	Bn		14	C	712 (M) ⁺
	43	Ms-D-Ser (Bn)	Bn		11	C	718 (M) ⁺


- 37 -




44	Ms-D-Ser (Bn)	Bn		31	D	729 (M+2) ⁺
45	Ms-D-Ser (Bn)	Bn		64	D	688
46	Ms-D-Ser (Bn)	Bn		144	D	660

*2:1 ratio of diastereomers. **Percent inhibition @ 10 μ M

5

Table 3

Inhibitory Activity of Branched-Chain α -Ketoamides

Ex. No.	D	Calpain IC ₅₀ nM	Prep. Method	MS (M+1)
10 47		26	A	645
48		43	A	645
49		32	A	646 (M+2)


- 38 -

50 $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{Ph}$ (100%) C 645

**

Compounds listed in Table 4 were prepared by the general methods A-G described above.

Table 4

Inhibitory Activity of α -Ketoamides

Ex. No.	W	R	Calpain IC ₅₀ nM	MS (M+1)
51	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(3-(2-NH}_2\text{-thiazol-4-yl)Ph)}$	29	729
52	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(5-(3-formylphenyl)thiophene-2-yl)}$	5	741
10 53	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHC(=N-CN)OPh}$	15	635
54	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(5-(3-(Me}_2\text{NCH}_2\text{)phenyl)thiophene-2-yl)}$	12	770
55	Ms-D-Ser (Bn)	3-Boc-NH-cyclohexane	42	645
56	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(5-(3-(morpholinoCH}_2\text{)phenyl)thiophene-2-yl)}$	18	812
57	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(4-(MorpholinoCH}_2\text{)Ph)}$	18	730
15 58	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(5-(3-(N-Me-piperazinyl-CH}_2\text{)phenyl)thiophene-2-yl)}$	21	825
59	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(5-(3-(HOCH}_2\text{)phenyl)thiophene-2-yl) (2 diast)}$	35	765 (M+Na)

	60	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{SO}_2\text{NHPH}$	47	631
	61	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{SO}_2\text{NH}(4\text{-CF}_3\text{Ph})$	32	699
	62	Ms-D-Ser (Bn)	$(\text{CH}_2)_3\text{SO}_2\text{NHPH}$	18	645
	63	Ms-D-Ser (Bn)	$(\text{CH}_2)_3\text{SO}_2\text{NH}(4\text{-CF}_3\text{Ph})$	23	713
5	64	Ms-D-Ser (Bn)	6-ketopiperidin-3-yl	(33)*	545
	65	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{N}(\text{Me})\text{SO}_2\text{-(5-(3-formylphenyl)thiophene-2-yl)}$	19	755
	66	Ms-D-Thr (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2(5\text{-pyrid-2-ylthiophene-2-yl})$	12	728
	67	Ms-D-Ser (Bn)	$\text{-N}(\text{Me})\text{SO}_2\text{-(5-isoxazol-3-yl-thiophene-2-yl)}$	21	718
	68	Ms-(D,L)- Phenylgly	$\text{CH}_2\text{CH}_2\text{NHSO}_2(5\text{-pyrid-2-ylthiophene-2-yl})$	21	670
10	69	Ms-(D,L)- Phenylgly	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{Ph}$	80	587
	70	Ms-D-Thr (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2(5\text{-(3-(morpholinoCH}_2\text{)phenyl)thiophene-2-yl})$ (Mixture of diastereomers)	23	826
	71	Ms-D-Phe	$\text{CH}_2\text{CH}_2\text{NHSO}_2(5\text{-pyrid-2-ylthiophene-2-yl})$	18	684
	72	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2(5\text{-(3-Fluorophenyl)thiophene-2-yl})$ (Mixture of diastereomers)	18	731
	73	Ms-D-Ser (Bn)	$(\text{CH}_2)_3\text{SO}_2\text{NHOCH}_3$	87	597 (M-1)
15	74	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(5-(3-Nitrophenyl)thiophene-2-yl)}$ (Mixture of diastereomers)	15	758
	75	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2\text{-(5-(3-Methylphenyl)thiophene-2-yl)}$ (Mixture of diastereomers)	36	727
	76	Ms-D-Ser (Bn)	$\text{CH}_2\text{CH}_2\text{NHSO}_2(5\text{-(3-(AcNH)phenyl)thiophene-2-yl})$ (Mixture of diastereomers)	11	792 (M+Na)

	77	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(CH ₃ CO)phenyl)thiophene-2-yl) (Mixture of diastereomers)	10	777 (M+Na)
	78	Ms-D-Ser (Bn)	1-(4-(Morpholinomethyl)benzenesulfonyl)piperidin-4-yl) (Mixture of diastereomers)	48	770
	79	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ ((4-(CH ₃ COPh)piperazin-1-yl)CH ₂ Ph) (Mixture of diastereomers)	11	847
	80	Ms-D-Ser (Bn)	(CH ₂) ₃ SO ₂ NH-morpholin-4-yl	169	652 (M-1)
5	81	Ms-D-Ser (Bn)	(CH ₂) ₃ SO ₂ -morpholin-4-yl	124	639
	82	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ -(5-(4-Methoxyphenyl)thiophene-2-yl) (Mixture of diastereomers)	13	765 (M+Na)
	83	Ms-D-Ser (Bn)	CH ₂ CH ₂ CH ₂ -Saccharin	48	657
	84	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ ((4-(PhCH ₂)piperazin-1-yl)CH ₂ Ph)	23	819
	85	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ ((4-(CH ₃ CO)piperazin-1-yl)CH ₂ Ph)	14	771
10	86	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ -(5-Me ₂ N-naphth-1-yl)	49	724
	87	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ -benzothiophene-2-yl	23	687
	88	Cbz-Leu-Leu	CH ₂ CH ₂ NHSO ₂ (5-pyrid-2-ylthiophene-2-yl)	33	819
	89	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ ((4-Pyrid-2-yl)piperazin-1-yl)CH ₂ Ph) (Mixture of diastereomers)	21	806
	90	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ -(5-(4-formylphenyl)thiophene-2-yl)	17	741
15	91	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (4-(2-(MeOCH ₂)Pyrrolidinyl)CH ₂)Ph)	19	758

	92	Ms-D-Ser (Bn)	(CH ₂) ₅ NHSO ₂ (5-pyrid-2-ylthiophene-2-yl) (Mixture of diastereomers)	12	756
	93	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(2-(morpholinoCH ₂)phenyl)thiophene-2-yl)	40	812
	94	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(4-(morpholinoCH ₂)phenyl)thiophene-2-yl)	22	812
	95	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(piperidinylCH ₂)phenyl)thiophene-2-yl)	30	810
5	96	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (2-acetamido-4-methylthiazol-5-yl)	23	709
	97	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (1-phenylsulfonylpiperidin-4-yl)	32	671
	98	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(2-formylphenyl)thiophene-2-yl)	24	741
	99	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ ((CH ₃ O)CH ₃ NCH ₂ Ph) (Mixture of diastereomers)	21	704
	100	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (4-ethylpiperazin-1-yl)CH ₂ Ph) (Mixture of diastereomers)	21	756
10	101	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(Et ₂ NCH ₂)phenyl)thiophene-2-yl)	22	798
	102	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(Cyclohexyl(Me)NCH ₂)phenyl)thiophene-2-yl)	36	838
	103	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(pyrrolidinylCH ₂)phenyl)thiophene-2-yl)	24	796
	104	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-cyanophenyl)thiophene-2-yl)	10	738
	105	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (4-(4-acetamidophenoxy)CH ₂ Ph)	14	816 (M+Na)
15	106	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(azetidiny)CH ₂)phenyl)thiophene-2-yl)	44	782


- 42 -

	107	Ms-D-Ser (Bn)	1-(5-pyridin-2-ylthiophene-2-yl- SO ₂)Piperidin-4-yl) (Mixture of diastereomers)	23	754
	108	Ms-D-Ser (Bn)	CONHCH ₂ CH ₂ NHSO ₂ (5-(3-(N-ethyl-N- methylaminomethyl)phenyl)thiophene- 2-yl)	10	784
	109	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(bis(2- methoxyethyl)aminomethyl)phenyl)thiophene-2- yl)	22	858
	110	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-cyanophenyl)thiophene-2- yl) (Mixture of diastereomers)	11	738
5	111	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (4-(3-pyrrolin-1-yl)CH ₂ Ph) (Mixture of diastereomers)	73	712
	112	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ ((4-(CH ₃ SO ₂)piperazin-1- yl)CH ₂ Ph)	37	807
	113	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ ((4-pyrimid-2-yl)piperazin-1- yl)CH ₂ Ph	24	807
	114	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3- (thiomorpholinoCH ₂)phenyl)thiophene-2-yl)	33	828
	115	Ms-D-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ (5-(3-(4- ketopiperidinyl)CH ₂)phenyl)thiophene-2-yl)	16	824
10	116	Ms-L-Ser (Bn)	CH ₂ CH ₂ NHSO ₂ Ph	100	631





















*Percent inhibition @ 0.1 μM

Ms = methylsulfonyl





Table 5

Inhibitory Activity of Achiral P₂ Mimetic α -Ketoamides

Ex. No.	Q	R	Calpain IC ₅₀ nM	Synthesis Method	MS (M+1) ⁺
117	Benzoyl		800	A	563
118	2,6-Dichlorobenzoyl		36	A	631, 633
119	2,6-Dichloro-3-methylbenzoyl		61	A	645
120	2,6-Difluorobenzoyl		20	A	599
121	2,4,6-Trifluorobenzoyl		85	A	618
122	2,3,4,5,6-Pentafluorobenzoyl		28	A	653
123	3,4-Methylenedioxybenzoyl		>1000	A	607
124	2,5-Dichlorobenzoyl		68	A	631, 633
125	2-Chloro-5-methoxybenzoyl		65	A	627
126	3,5-bis(trifluoromethyl)benzoyl		600	A	699
127	2,6-Dimethylbenzoyl		178	A	591

	128	2,6-Dichloronicotinoyl		80	A	634, 636
	129	2,6-Dichlorobenzoyl		21	A	658, 660
	130	2,6-Dichlorobenzoyl		20	A	687, 689
	131	2,6-Dichlorobenzoyl		29	A	729, 731
5	132	2,6-Dichlorobenzoyl		83	A	723, 725
	133	2,6-Dichlorobenzoyl		11	A	655, 657
	134	2,6-Dichlorobenzoyl		100	A	688, 690
	135	2,6-Difluorobenzoyl		22	A	645 (M+Na)
	136	2,6-Difluorobenzoyl		62	A	611 (M+Na)
10	137	2,6-Diethylbenzoyl		145	A	631 (M+Na)
	138	2,6-Dimethoxybenzoyl		4000	A	613
	139	2-Isopropylbenzoyl		168	A	595
	140	2-Chloro-6-fluorobenzoyl		58	A	605
	141	2-Fluoro-6-trifluoromethylbenzoyl		58	A	639
15	142	2,3,4,5,6-Pentafluorobenzoyl		32	A	643
	143	2-Methylpropanoyl		1500	A	529
	144	3-Methylbutanoyl		590	A	543
	145	4-Methylpentanoyl		29	A	557
	146	3-Cyclopentylpropanoyl		1000	A	583
20	147	E-3-Hexenoyl		1000	A	555

- 45 -

148	4-Phenylpentanoyl		87	A	641 (M+Na)
149	4-Phenylbutanoyl		1500	A	627 (M+Na)
150	4-Methylpentanoyl		15	A	603 (M+Na)
151	3-Cyclopentylpropanoyl		420	A	607


5 As those skilled in the art will appreciate, numerous changes and modifications may be made to the preferred embodiments of the invention without departing from the spirit of the invention. It is intended that all such variations fall within the scope of the invention.

10 It is intended that each of the patents, applications, and printed publications mentioned in this specification be hereby incorporated by reference in their entirety.

- 46 -

What is claimed is:

1. A compound having the Formula I:



I

5 wherein:

Q has the formula G-B-(CHR⁴)_v where R⁴ is independently H or alkyl having from 1 to 4 carbons;

v is 0, 1, or 2;

B is selected from the group consisting of C(=O),
 10 OC(=O), S(=O)_m, CH₂, a bond, NR⁵C(=O), S(=O)_m-A-C(=O), and
 C(=O)-A-C(=O), where R⁵ is H or lower alkyl;

m is 0, 1, or 2;

A is lower alkylene or cycloalkylene,
 optionally substituted with one or more halogen atoms,
 15 aryl, or heteroaryl groups;

M is a carbon atom;

G is selected from the group consisting of H, a
 blocking group, lower alkyl, lower alkenyl, aryl having
 from about 6 to about 14 carbons, heterocyclyl having from
 20 about 5 to about 14 ring atoms, heterocycloalkyl having
 from about 5 to about 14 ring atoms, arylalkyl having from
 about 7 to about 15 carbons, heteroarylalkyl, and
 arylheteroalkyl wherein the aryl portion can be unfused or
 fused with the heteroalkyl ring, said alkyl, aryl,
 25 heterocyclyl, heterocycloalkyl, arylalkyl, heteroarylalkyl,
 and arylheteroalkyl groups being optionally substituted
 with one or more J groups;

J is selected from the group consisting of halogen,
 CN, nitro, lower alkyl, cycloalkyl, heterocycloalkyl,
 30 heteroalkyl, halogenated alkyl, aryloxyalkyl, alkylthio,
 alkylsulfonyl, aryl, heteroaryl, arylalkyl, arylalkyloxy,

- 47 -

arylsulfonyl, heteroarylsulfonyl, alkoxycarbonyl, alkoxyalkyl, acyl, alkoxy, hydroxy, carboxy, hydroxyalkyl, amino, alkylamino, and aminoalkyl, said amino group or said amino group of said aminoalkyl or alkylamino group being
5 optionally substituted with an acyl group, an alkoxy group, or with 1 to 3 aryl, lower alkyl, cycloalkyl, or alkoxyalkyl groups; and said aryl, heteroaryl, heterocycloalkyl, and heteroalkyl groups being further optionally substituted by a J group;

10 each Aaa is independently an amino acid which optionally contains one or more blocking groups;

n is 0, 1, 2, or 3;

R¹ and R² are independently selected from the group consisting of H, alkyl having from one to about 6 carbons,
15 arylalkyl having from about 7 to about 15 carbons, heteroalkyl in which the ring contains from about 5 to about 14 ring atoms, heteroarylalkyl in which the heteroaryl ring contains from about 5 to about 14 ring atoms, alkoxyalkyl, a side chain of a naturally occurring
20 amino acid in the R or S configuration, and (CH₂)_pNH-L, said alkyl, arylalkyl, heteroalkyl, heteroarylalkyl, and alkoxyalkyl groups being optionally substituted with one or more J groups;

p is 0, 1, 2, or 3;

25 L is selected from the group consisting of alkoxycarbonyl having from 2 to about 7 carbons, arylalkoxycarbonyl in which the arylalkoxy group contains about 7 to about 15 carbons, and S(=O)₂R⁶;

R⁶ is selected from the group consisting of
30 lower alkyl, and aryl having from about 6 to about 14 carbons;

R³ is selected from the group consisting of H, alkyl having from one to about 6 carbons, arylalkyl having from about 7 to about 15 carbons, heteroalkyl in which the ring
35 contains from about 5 to about 14 ring atoms,


- 48 -

heteroarylalkyl in which the heteroaryl ring contains from about 5 to about 14 ring atoms, alkoxyalkyl, a side chain of a naturally occurring amino acid in the R or S configuration, $(CH_2)_pNH-L$, $C(=O)R^7$, $S(=O)_2R^7$, a blocking group, and when combined with the carbon atom to which R^1 is attached an alkylene group having from 2 to 5 carbons, said alkylene group being optionally substituted with a group selected from the group consisting of aryl, azide, CN, a protected amino group, and OSO_2 -aryl, said alkyl, arylalkyl, heteroalkyl, heteroarylalkyl, and alkoxyalkyl groups being optionally substituted with one or more J groups;

R^7 is selected from the group consisting of aryl having from about 6 to about 14 carbons, heteroaryl having from about 5 to about 14 ring atoms, arylalkyl having from about 7 to about 15 carbons, alkyl having from 1 to about 10 carbons, said aryl, heteroaryl, arylalkyl and alkyl groups being optionally substituted with one or more J groups, heteroalkyl having from 2 to about 7 carbons, alkoxy having from about 1 to about 10 carbons, and amino optionally substituted with 1 or more alkyl groups;

q is 0 or 1;

Z is selected from the group consisting of $C(=O)C(=O)NH-X-A^1-K$ and




25 X is a bond or -O-;

A^1 is the same as A;

K is selected from the group consisting of $N(R^{10})Y$,

- 49 -



D is a fused aryl or heteroaryl group;

R^{11} is selected from the group consisting of alkoxy, aryloxy, and NHR^{12} ;

5 R^{12} is selected from the group consisting of H, alkyl, aryl, and heteroaryl, said alkyl, aryl or heteroaryl groups being optionally substituted with one or more J groups;

Y is selected from the group consisting of SO_2R^8 ,
 10 $\text{C}(=\text{O})\text{NHR}^9$, $\text{C}(=\text{S})\text{NHR}^9$, $\text{C}(=\text{NCN})\text{R}^{11}$, $\text{C}(=\text{NC}(=\text{O})\text{NHR}^{10})\text{R}^{11}$, and CO_2R^8 ;

R^8 is selected from the group consisting of alkyl, alkoxy, aryl, and heterocyclyl, said alkyl, alkoxy, aryl, or heterocyclyl groups being optionally substituted
 15 with one or more J groups;

R^9 is selected from the group consisting of H, alkyl, aryl, and heteroaryl, said alkyl, aryl, or heteroaryl groups being optionally substituted with one or more J groups;

20 or an R^9 alkyl group may be combined with an A^1 alkylene group to form a N-containing heterocyclic 5- or 6-membered ring;

R^{10} is selected from the group consisting of H and lower alkyl;

25 or in the moiety $\text{SO}_2\text{N}(\text{R}^8)\text{R}^{10}$, R^8 and R^{10} may be combined together with the N atom to which they are attached to form a N-containing heterocyclic 5- or 6-membered ring;

or where A^1 is an alkylene group, and K is $\text{N}(\text{R}^{10})\text{Y}$
 30 wherein R^{10} is alkyl, said R^{10} alkyl group may be combined

- 50 -

with said A¹ alkylene group to form a N-containing heterocyclic 5- or 6- membered ring;
or a pharmaceutically acceptable salt thereof.

2. The compound of claim 1 wherein n and v are each
5 0, q is 1, B is a bond, and G is H.

3. The compound of claim 1 wherein R¹ is the sidechain of a naturally occurring amino acid.

4. The compound of claim 1 wherein R³ is
-S(=O)₂R⁷.

10 5. The compound of claim 1 wherein R² is benzyl or alkoxyalkyl.

6. The compound of claim 1 wherein X is a bond, and Y is SO₂R⁸.

7. The compound of claim 1 wherein A¹ is
15 -CH₂-CH₂-, -CH₂-CH(CH₃)-, or -(CH₃)CH-CH₂-.

8. The compound of claim 1 wherein R¹ is a serine sidechain, which is optionally capped with a benzyl group.

9. The compound of claim 8 wherein M is a carbon
20 atom in the D configuration.

10. The compound of claim 1 wherein R² is benzyl, R⁷ is methyl, and R⁸ is substituted phenyl, unsubstituted phenyl, substituted heteroaryl, or unsubstituted heteroaryl.

25 11. The compound of claim 1 wherein R⁸ is aryl, aryl substituted with amino, aryl substituted with

- 51 -

heterocyclomethyl, heteroaryl, alkyl substituted with heteroaryl, or heteroaryl substituted with alkylthio, haloalkyl, alkyl, phenylsulfonyl, halogen, aminophenyl, amino, or dialkylaminoalkyl.

5 12. The compound of claim 1 wherein n and v are each 0, q is 1, R¹ is the side chain of an amino acid in the D- or L-configuration, R³ is S(=O)₂R⁷, G is H, B is a bond, R² is benzyl or alkoxyalkyl, X is a bond, and Y is SO₂R⁸.


10 13. The compound of claim 1 wherein A¹ is CH₂CH₂, CH₂CH(CH₃), or (CH₃)CHCH₂.

14. The compound of claim 1 wherein R¹ is a serine side chain in the D-configuration in which the hydroxyl group is capped with benzyl, R² is benzyl, R⁷ is methyl, and
15 R⁸ is substituted or unsubstituted phenyl or substituted or unsubstituted heteroaryl.

15. The compound of claim 1 wherein R₁-R₄, B, G, Aaa, X, A¹, Y, n, q and v are selected in accordance with Tables
20 2 and 3.

16. The compound of claim 1 wherein R₁-R₄, B, G, Aaa, X, A¹, Y, n, q and v are each independently selected from the group of substituents shown in Tables 2 and 3.

17. The compound of claim 1 having the Formula:




wherein:


W, R₂ and R are independently selected from the group of substituents shown in Table 2.

18. The compound of claim 17 wherein W, R₂ and R are selected in accordance with Table 2.


19. The compound of claim 1 having the Formula:



5 wherein D is $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)\text{SO}_2\text{Ph}$ or has one of the formulas:




20. The compound of claim 1 having the formula:



wherein W and R are independently selected from the
10 group of substituents shown in Table 4.

21. The compound of claim 20 wherein W and R are selected in accordance with Table 4.

22. The compound of claim 1 having the Formula:



- 53 -

wherein Q and R are independently selected from the group of substituents shown in Table 5.

23. The compound of claim 22 wherein Q and R are selected in accordance with Table 5.

5 24. The compound of claim 1 wherein n, v and q are each 0; B is (C=O); and G is phenyl or lower alkyl, said phenyl or lower alkyl groups being optionally substituted with one or more J groups.

25. A composition for inhibiting a serine protease or
10 a cysteine protease comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

26. A method for inhibiting a serine protease or a cysteine protease comprising contacting a protease selected from the group consisting of serine proteases and cysteine
15 proteases with an inhibitory amount of a compound of claim 1.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/21055

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 38/05, 38/06; C07K 5/00

US CL : 514/18-19; 530/331

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/18-19; 530/331

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN STRUCTURE & BIBLIOGRAPHIC LIBRARY SEARCHES

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	Chemical Abstract Vol. 129, April 1998, Abstract No. 28214, BIHOVSKY et al., WO 9821186 A1, 22 MAY 1998, see chemical structures of RN 208051-55-2 and 208052-81-7.	1-11, 13-15 and 25-26

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
28 DECEMBER 1998

Date of mailing of the international search report
26 JAN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

BENNETT CELSA

Telephone No. (703) 308-0196